



BRIGHT FUTURES HANDOUT ► PARENT

2 MONTH VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.

✓ HOW YOUR FAMILY IS DOING

- If you are worried about your living or food situation, talk with us. Community agencies and programs such as WIC and SNAP can also provide information and assistance.
- Find ways to spend time with your partner. Keep in touch with family and friends.
- Find safe, loving child care for your baby. You can ask us for help.
- Know that it is normal to feel sad about leaving your baby with a caregiver or putting him into child care.

✓ HOW YOU ARE FEELING

- Take care of yourself so you have the energy to care for your baby.
- Talk with me or call for help if you feel sad or very tired for more than a few days.
- Find small but safe ways for your other children to help with the baby, such as bringing you things you need or holding the baby's hand.
- Spend special time with each child reading, talking, and doing things together.

✓ FEEDING YOUR BABY

- Feed your baby only breast milk or iron-fortified formula until she is about 6 months old.
- Avoid feeding your baby solid foods, juice, and water until she is about 6 months old.
- Feed your baby when you see signs of hunger. Look for her to
 - Put her hand to her mouth.
 - Suck, root, and fuss.
- Stop feeding when you see signs your baby is full. You can tell when she
 - Turns away
 - Closes her mouth
 - Relaxes her arms and hands
- Burp your baby during natural feeding breaks.

If Breastfeeding

- Feed your baby on demand. Expect to breastfeed 8 to 12 times in 24 hours.
- Give your baby vitamin D drops (400 IU a day).
- Continue to take your prenatal vitamin with iron.
- Eat a healthy diet.
- Plan for pumping and storing breast milk. Let us know if you need help.
 - If you pump, be sure to store your milk properly so it stays safe for your baby. If you have questions, ask us.

If Formula Feeding

- Feed your baby on demand. Expect her to eat about 6 to 8 times each day, or 26 to 28 oz of formula per day.
- Make sure to prepare, heat, and store the formula safely. If you need help, ask us.
- Hold your baby so you can look at each other when you feed her.
- Always hold the bottle. Never prop it.

✓ YOUR GROWING BABY

- Have simple routines each day for bathing, feeding, sleeping, and playing.
- Hold, talk to, cuddle, read to, sing to, and play often with your baby. This helps you connect with and relate to your baby.
- Learn what your baby does and does not like.
- Develop a schedule for naps and bedtime. Put him to bed awake but drowsy so he learns to fall asleep on his own.
- Don't have a TV on in the background or use a TV or other digital media to calm your baby.
- Put your baby on his tummy for short periods of playtime. Don't leave him alone during tummy time or allow him to sleep on his tummy.
- Notice what helps calm your baby, such as a pacifier, his fingers, or his thumb. Stroking, talking, rocking, or going for walks may also work.
- *Never hit or shake your baby.*

Helpful Resources:

Information About Car Safety Seats: www.safercar.gov/parents | Toll-free Auto Safety Hotline: 888-327-4236

2 MONTH VISIT—PARENT



SAFETY

- Use a rear-facing-only car safety seat in the back seat of all vehicles.
- Never put your baby in the front seat of a vehicle that has a passenger airbag.
- Your baby's safety depends on you. Always wear your lap and shoulder seat belt. Never drive after drinking alcohol or using drugs. Never text or use a cell phone while driving.
- Always put your baby to sleep on her back in her own crib, not your bed.
 - Your baby should sleep in your room until she is at least 6 months old.
 - Make sure your baby's crib or sleep surface meets the most recent safety guidelines.
- If you choose to use a mesh playpen, get one made after February 28, 2013.
- Swaddling should not be used after 2 months of age.
- Prevent scalds or burns. Don't drink hot liquids while holding your baby.
- Prevent tap water burns. Set the water heater so the temperature at the faucet is at or below 120°F /49°C.
- Keep a hand on your baby when dressing or changing her on a changing table, couch, or bed.
- Never leave your baby alone in bathwater, even in a bath seat or ring.

WHAT TO EXPECT AT YOUR BABY'S 4 MONTH VISIT

We will talk about

- Caring for your baby, your family, and yourself
- Creating routines and spending time with your baby
- Keeping teeth healthy
- Feeding your baby
- Keeping your baby safe at home and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th Edition

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®




The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.



BIRTH TO 6 MONTHS

Safety for Your Child

Did you know that hundreds of children younger than 1 year die every year in the United States because of injuries—most of which could be prevented?

Often, injuries happen because parents are not aware of what their children can do. Children *learn quickly*, and before you know it, your child will be *wiggling* off a bed or *reaching* for your cup of hot coffee.

Car Injuries

Car crashes are a great threat to your child's life and health. Most injuries and deaths from car crashes **can be prevented** by the use of car safety seats. Your child, besides being much safer in a car safety seat, will behave better, so you can pay attention to your driving. Make your newborn's first ride home from the hospital a safe one—in a car safety seat. Your infant should ride in the back seat in a rear-facing car safety seat.

Make certain that your baby's car safety seat is installed correctly. Read and follow the instructions that come with the car safety seat and the sections in the owners' manual of your car on using car safety seats correctly. Use the car safety seat EVERY time your child is in the car.

NEVER put an infant in the front seat of a car with a passenger airbag.



Falls

Babies *wiggle* and *move* and *push* against things with their feet soon after they are born. Even these very first movements can result in a fall. As your baby grows and is able to roll over, he or she may fall off of things unless protected. **Do not leave your baby alone** on changing tables, beds, sofas, or chairs. **Put your baby in a safe place** such as a crib or playpen when you cannot hold him or her.

Your baby may be able to crawl as early as 6 months. **Use gates on stairways and close doors** to keep your baby out of rooms where he or she might get hurt. **Install operable window guards** on all windows above the first floor.

Do not use a baby walker. Your baby may tip the walker over, fall out of it, or fall down stairs and seriously injure his or her head. Baby walkers let children get to places where they can pull heavy objects or hot food on themselves. A better choice is a stationary activity center with no wheels.

If your child has a serious fall or does not act normally after a fall, call your doctor.



(over)

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



Burns

At 3 to 5 months, babies will wave their fists and grab at things. **NEVER carry your baby and hot liquids, such as coffee, or foods at the same time.** You can't handle both. Your baby can get burned! To protect your child from tap water scalds, the hottest temperature at the faucet should be no more than 120°F. In many cases you can adjust your water heater.

If your baby gets burned, immediately put the burned area in cold water. Keep the burned area in cold water for a few minutes to cool it off. Then cover the burn loosely with a dry bandage or clean cloth and call your doctor.


To protect your baby from house fires, be sure you have a working smoke alarm on every level of your home, especially in furnace and sleeping areas. Test the alarms every month. It is best to use smoke alarms that use long-life batteries, but if you do not, change the batteries at least once a year.

Choking and Suffocation

Babies explore their environment by putting anything and everything into their mouths. **NEVER leave small objects in your baby's reach, even for a moment.** NEVER feed your baby hard pieces of food such as chunks of raw carrots, apples, hot dogs, grapes, peanuts, and popcorn. Cut all the foods you feed your baby into thin pieces to prevent choking. **Be prepared if your baby starts to choke. Ask your doctor to recommend the steps you need to know. Learn how to save the life of a choking child.**

To prevent possible suffocation and reduce the risk of sudden infant death syndrome (SIDS), **your baby should always sleep on his or her back. Your baby should have his or her own crib or bassinet with no pillows, stuffed toys, bumpers, or loose bedding. NEVER put your baby on a waterbed, beanbag, or anything that is soft enough to cover the face and block air to the nose and mouth.**

Plastic wrappers and bags form a tight seal if placed over the mouth and nose and may suffocate your child. Keep them away from your baby.



From Your Doctor

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.